

POSTER PRESENTATION

Open Access

Expression of TAAs in glioblastoma and expansion of anti-TAA -reactive T cells

Ernest Dodoo¹, Liu zhenjiang², Bartek Jiri², Oscar Persson², Qingda Meng², Thomas Poirer², Lalit Rane², Christopher Illies², Julia Karbach³, Elke Jäger³, Markus Maeurer^{2*}

From Society for Immunotherapy of Cancer 29th Annual Meeting
National Harbor, MD, USA. 6-9 November 2014

Purpose

Active cellular therapy (ACT) using *ex-vivo* expanded T cells from patients with cancer, obtained by apheresis, can represent a viable source for anti-cancer directed cellular therapy. TAAs expressed in glioblastoma may represent attractive targets for i) CARs, ii) transgenic T cells targeting nominal tumor antigens (e.g. NY-ESO-1) or iii) T cells enriched for TAA after *ex vivo* expansion.

Methods

Fresh blood samples were obtained from 50 patients with tumors of the central nervous system and tested for anti-TAA reactivity. T cells were expanded without cytokines, with IL-2 and IL-7, or with IL-2, IL-15 and IL-21 and tested for CD4/8 expansion by flow cytometry and for IFN-gamma production by ELISA. PBMCs were expanded using IL-2/15/21 and peptides (15mers) covering TAAs (surviving or NY-ESO-1). The T cell phenotype (CD3, CD4, CD8, CD45RA and CCR7) was determined by flow cytometry and TAA-reactive T cells were identified by intracellular cytokine staining (IL-2, TNF, IFN and IL-17). TAA-specific IgG in serum was detected by a quantitative ELISA in patients with glioblastoma and in age- and sex-matched healthy donors. FGFRVIII was determined by RT-PCR and protein expression of survivin and NY-ESO-1 was evaluated by immunohistochemistry and graded using a scale from 1+ to 4+ along with pattern analysis of TAA expression.

Results

We could detect IFN-gamma responses in 25% blood samples for NY-ESO-1 and 30% for survivin and antigen-specific CD8 / CD4+ T cell proliferation. Cellular

responses could be augmented by adding cytokines, i.e. IL-2 and IL-7 favored CD4+ T cell proliferation, IL-2, IL-15 and IL-21 favored CD8+ T cell proliferation. TAAs-reactive T cells could be successfully expanded *ex vivo* and exhibited TAA-specific production of IFN-gamma and TNFalpha and a CD8+CD45RA-CCR7+ phenotype. 38/50 specimens expressed NY-ESO-1, yet only 3/50 showed a strong, universal (4+) NY-ESO-1 protein expression pattern, 12/50 cancer lesions exhibited a strong (4+) staining for survivin defined by immunohistochemistry. 25 % of glioblastoma tested positive for the FGFRVIII.

Conclusion

A TAA-specific WBA (whole blood assay) can be used to gauge the potential for expansion of TAA-reactive T cells in peripheral blood from patients with glioblastoma. TAA-reactive T cells can be successfully expanded from patients with glioblastoma in IL-2, IL-15 and IL-21, they exhibit a central memory phenotype and produce a Th1 cytokine cytokine pattern. NY-ESO-1 expression in glioblastoma represents a viable target for anti-NY-ESO-1 directed T cells.

Consent

Written informed consent was obtained from the patient for publication of this abstract and any accompanying images. A copy of the written consent is available for review by the Editor of this journal.

Authors' details

¹Dept. of Neurosurgery, Karolinska University Hospital, Stockholm, Sweden.

²Karolinska Institutet, Stockholm, Sweden. ³Krankenhaus Nordwest, Frankfurt/M., Germany.

²Karolinska Institutet, Stockholm, Sweden

Full list of author information is available at the end of the article

Published: 6 November 2014

doi:10.1186/2051-1426-2-S3-P24

Cite this article as: Dodoo et al.: Expression of TAAs in glioblastoma and expansion of anti-TAA -reactive T cells. *Journal for ImmunoTherapy of Cancer* 2014 **2**(Suppl 3):P24.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

